

Photonic Stimulation (Photobiomodulation) - Clinical and Experimental Applications of NIR-LED

KRISTINA D. DESMET, B.S.,¹ DAVID A. PAZ, B.S.,² JESSE J. CORRY, M.D.,² JANIS T. EELLS, Ph.D.,¹ MARGARET T.T. WONG-RILEY, Ph.D.,³ MICHELE M. HENRY, B.S.,⁴ ELLEN V. BUCHMANN, B.S.,² MARY P. CONNELLY, B.S.,² JULIAV. DOVI, Ph.D.,² HUAN LING LIANG, M.D.,³ DIANE S. HENSHEL, Ph.D.,⁵ RONNIE L. YEAGER, M.S.,⁵ DEBORAH S. MILLSAP, M.S.,⁵ JINHWAN LIM, M.S.,⁵ LISA J. GOULD, M.D., Ph.D.,⁶ RINA DAS, Ph.D.,⁷ MARTI JETT, Ph.D.,⁷ BRIAN D. HODGSON, D.D.S.,⁸ DAVID MARGOLIS, M.D.,⁹ and HARRY T. WHELAN, M.D.²

ABSTRACT: This review presents current research on the use of far-red to near-infrared (NIR) light treatment in various in vitro and in vivo models. Low-intensity light therapy, commonly referred to as "photobiomodulation," also called "photonic stimulation," uses light in the far-red to near-infrared region of the spectrum (630–1000 nm) and modulates numerous cellular functions. Positive effects of NIR-light-emitting diode (LED) light treatment include acceleration of wound healing, improved recovery from ischemic injury of the heart, and attenuated degeneration of injured optic nerves by improving mitochondrial energy metabolism and production. Various in vitro and in vivo models of mitochondrial dysfunction were treated with a variety of wavelengths of NIR-LED light. These studies were performed to determine the effect of NIR-LED light treatment on physiologic and pathologic processes. NIR-LED light treatment stimulates the photoacceptor cytochrome c oxidase, resulting in increased energy metabolism and reduction. NIR-LED light treatment accelerates wound healing in ischemic rat and murine diabetic wound healing models, attenuates the retinotoxic effects of methanol-derived formic acid in rat models, and attenuates the developmental toxicity of dioxin in chicken embryos. Furthermore, NIR-LED light treatment prevents the development of oral mucositis in pediatric bone marrow transplant patients. The experimental results demonstrate that NIR-LED light treatment stimulates mitochondrial oxidative metabolism in vitro, and accelerates cell and tissue repair in vivo. NIR-LED light represents a novel, noninvasive, therapeutic intervention for the treatment of numerous diseases linked to mitochondrial dysfunction.

CONCLUSION:

Experimental results demonstrate that NIR-LED light treatment stimulates mitochondrial oxidative metabolism in vitro, and accelerates cell and tissue repair in vivo. NIR-LED light represents a noninvasive, therapeutic intervention for the treatment of numerous diseases linked to mitochondrial dysfunction, including age-related macular degeneration, Leber's hereditary optic neuropathy, and Parkinson's and Alzheimer's disease.

1. Department of Clinical Laboratory Sciences, University of Wisconsin–Milwaukee, Milwaukee, Wisconsin.
2. Department of Neurology, Medical College of Wisconsin, Milwaukee, Wisconsin.
3. Department of Cell Biology, Neurobiology, and Anatomy, Medical College of Wisconsin, Milwaukee, Wisconsin.
4. Department of Ophthalmology, Medical College of Wisconsin, Milwaukee, Wisconsin.
5. School of Public and Environmental Affairs, Indiana University, Bloomington, Indiana.
6. Department of Plastic Surgery, Medical College of Wisconsin, Milwaukee, Wisconsin.
7. Department of Molecular Pathology, Walter Reed Army Institute of Research, Silver Spring, Maryland.
8. Department of Dentistry, Children's Hospital of Wisconsin, Milwaukee, Wisconsin.
9. Department of Hematology and Oncology, Medical College of Wisconsin, Milwaukee, Wisconsin.